

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
1 March 2001 (01.03.2001)

PCT

(10) International Publication Number
WO 01/14361 A1

- (51) International Patent Classification⁷: C07D 311/82, Longfellow Road, Natick, MA 01760 (US). CRESPI, Charles, L.; 4 Evans Road, Marblehead, MA 01945 (US).
C12Q 1/00
- (21) International Application Number: PCT/US00/21894 (74) Agent: PLUMER, Elizabeth, R.; Wolf, Greenfield & Sacks, P.C., 600 Atlantic Avenue, Boston, MA 02210 (US).
- (22) International Filing Date: 10 August 2000 (10.08.2000) (81) Designated State (national): JP.
- (25) Filing Language: English (84) Designated States (regional): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
- (26) Publication Language: English
- (30) Priority Data:
60/149,762 19 August 1999 (19.08.1999) US
60/150,044 20 August 1999 (20.08.1999) US
Published:
— With international search report.
— Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.
- (71) Applicant: GENTEST CORPORATION [US/US]; 6 Henshaw Street, Woburn, MA 01801 (US).
- (72) Inventors: MILLER, Vaughn, P.; 36 Marathon Street, Arlington, MA 02474 (US). STRESSER, David; 31

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

BEST AVAILABLE COPY



WO 01/14361 A1

(54) Title: THE USE OF FLUORESCIN ARYL ETHERS IN HIGH THROUGHPUT CYTOCHROME P450 INHIBITION ASSAYS

(57) Abstract: Novel fluorescent substrates of human cytochrome P450 enzymes are provided. Also provided are methods for their manufacture and use. These substrates are useful in assessing cytochrome P450 enzyme activity and in selecting compounds which inhibit cytochrome P450 enzyme activity and, in particular, for identifying potential adverse drug interactions which are mediated by inhibition of cytochrome P450 enzyme activity.

**The Use of Fluorescein Aryl Ethers in High Throughput
Cytochrome P450 Inhibition Assays**

Field of Invention

5 This invention relates to the field of drug and xenobiotic metabolism. The invention includes novel cytochrome P450 fluorescent probe substrates and reaction products, methods for their preparation and their use as assay reagents.

Background of the Invention

10 Cytochromes P450 (CYP) are the principal enzymes for the oxidative metabolism of many drugs, procarcinogens, promutagens, and environmental pollutants. Cytochrome P450 is a heme-containing, membrane-bound, multienzyme system that is present in many tissues *in vivo* but is present at the highest level in liver. In human liver, it is estimated that there are 15-20 different xenobiotic-metabolizing cytochrome P450 forms. A standard nomenclature based
15 on relatedness of amino acid sequences has been developed. A relatively limited subset of these enzymes, CYP1A2, CYP2C8, CYP2C9, CYP2C19, CYP2D6 and CYP3A4 appear to be most commonly responsible for the metabolism of drugs and associated drug-drug interactions (*Spatzenegger and Jaeger, 1995*). The relative importance of this subset of enzymes is due to:
20 the mass abundance of these enzymes (e.g. CYP3A4 is the most abundant P450 in human liver at ~30% of total P450), the preference of these enzymes to bind and/or metabolize chemical
structures commonly found in drugs (e.g. CYP2D6 preferentially binds and metabolizes drugs with basic amine functionalities), enzyme polymorphisms (such as CYP3A4 and CYP2C19), and enzyme regulation in response to environmental chemicals some (such as CYP1A2 and CYP3A4). Competition for metabolism by a particular cytochrome P450 form is a principal
25 mechanism of some clinically significant drug-drug interactions.

Identification of the enzymes responsible for metabolism is becoming an important aspect of drug development. Such identifications consider both the metabolism of the new drug as well as inhibition by the new drug. The identification of enzymes involved in metabolism of the new drug allows prediction, based on knowledge of the ability of
30 coadministered drugs to inhibit the same enzymes, of which coadministered drugs may inhibit the metabolism of the new drug. This information can also be used to predict individual

variability based on known metabolic polymorphisms. The identification of the enzymes most sensitive to inhibition by the new drug allows prediction, based on knowledge of which coadministered drugs are metabolized by the same enzyme, of which coadministered drug's metabolism may be inhibited by the new drug. Obtaining information for a series of drug candidates early in the drug discovery process can assist in the choice of the best drug candidate for further development.

Most cytochrome P450 assays have focused on the metabolism of drug molecules or drug candidates. While these chemicals are effective in assessing cytochrome P450 activity and inhibition, they are not amenable to high throughput screening assay technology (they require time consuming separation of enzyme reaction products using HPLC). Also, most of these substrates do not have the necessary fluorescent properties that make the substrate useful for in situ fluorescent plate analysis.

Sensitive fluorescent assays have been developed for human CYP1A1 and CYP1A2 enzymes. Members of the CYP1A family preferentially oxidize planar aromatic molecules not unlike the structure of common fluorescent chromophores. Microtiter plate based assays have been developed based on the O-dealkylation of alkyl ethers of resorufin (Donato, M. T. et al., *Anal. Biochem.* 213, 29-33 (1993); Kennedy, S. W. and S. P. Jones, *Anal. Biochem.*, 222, 217-223 (1994)) and coumarin (C. L. Crespi et al. *Anal Biochem.* 248, 188-190, (1997)). An intact cell assay using an alkyl ether of fluorescein (Quan, T. et al., *Carcinogenesis* 15, 1827-32, (1994)) has been described.

The development of fluorescent assays for other cytochromes P450 has been more challenging. We have described microtiter plate based assays for CYP2D6 based on alkylcoumarin derivatives (C. L. Crespi et al. *Anal Biochem.* 248, 188-190, (1997); Miller, V. P. and C. L. Crespi, U S patent application 09/352,576, entitled "Novel CYP2D Fluorescent Assay Reagents," filed July 12, 1999.) We have also reported the use of commercial alkylcoumarins for the fluorescent assay of CYP2C9 and CYP2C19. However, the sensitivity (signal to noise ratio) of these CYP2C assays is limited. New substrates which increase the assay signal and minimize the required amount of enzyme reagent would be useful. No fluorescent substrate has been reported for the important drug metabolizing enzyme CYP2C8. CYP2C8 is the primary cytochrome P450 enzyme responsible for the metabolism of the anticancer drug paclitaxel (TAXOLTM).

Among the different cytochromes P450 expressed in the liver, CYP3A4 is the most abundant. Important classes of substrates for this enzyme include steroids, macrolide antibiotics, antivirals, and polycyclic aromatic hydrocarbons. As the majority of commercially available drugs are metabolized by CYP3A4, the importance of screening for inhibition of this enzyme is essential in drug development. We have previously reported the use of the commercially available 7-benzyloxyresorufin (BzRes) as a fluorescent substrate for assessing CYP3A4 activity in a high throughput mode (Crespi et al. *Anal Biochem.* 248, 188-190, (1997)). We and others have demonstrated that the inhibition potential (e.g. IC_{50} value) or activation potential for most compounds varies dramatically depending on the probe substrate used (KE Thummel and GR Wilkinson, *Ann. Rev. Pharmacol. Toxicol.* 38:389-430 (1998)). Additional fluorescent probe substrates, which differ significantly in structure and chemical properties from the current substrates, are needed to gain a better understanding of the potential for test compounds to inhibit CYP3A4.

Fluorescein ethers have been reported to be useful fluorescent cytochrome P450 assay reagents. A. G. Miller (A. G. Miller, *Anal. Biochem.* 133, 46-57 (1983)) described a series of ethylated fluoresceins. Ethoxyfluorescein ethyl ether, reportedly was the most useful for the flow cytometric analysis and sorting of intact, viable rat cells based on cytochrome P450 activity. Other laboratories have also reported using the diethoxy- or a dimethoxyfluorescein derivative for the same purpose with rat or porcine cells (White et al., *Biochem. J.* 247, 23-28 (1987); Pan et al, *Artif. Organs* 20, 1173-1180 (1996); Anderson et al., *Int. J. Artif. Organs* 21, 360-364, (1998)). More recently, ethoxyfluorescein ethyl ester has been reported to monitor the activity of human skin fibroblasts transfected with human CYP1A1 cDNA (Quan, T. et al., *Carcinogenesis* 15, 1827-32, (1994)).

Aryl ether derivatives of fluorescein have not been reported as substrates for cytochrome P450 enzymes. A benzyl ether derivative of fluorescein, benzyloxyfluorescein benzyl ester, has been reported as an intermediate in the synthesis of fluorescein labeled polymers (Hargreaves and Webber, *Can. J. Chem.* 63, 1320-1327 (1985)), chiral prodyes (Tadic and Brossi, *Heterocycles* 31, 1975-1982, (1990)), and dye precursors (Dombrowski Jr. et al. US patent 5,196,297).


Summary of the Invention

The present invention relates to novel fluorescent substrates of cytochrome P450 enzymes. These substrates are useful in assessing cytochrome P450 enzyme activity and in selecting compounds which inhibit cytochrome P450 enzyme activity. Accordingly, the compounds and methods of the invention are useful for identifying potential adverse drug interactions which are mediated by inhibition of cytochrome P450 enzyme activity.

The compounds of the invention are substrates that are characterized in having properties which permit the sensitive quantitation of CYP2C8, CYP2C9, CYP2C19, and CYP3A activity using in situ fluorescence analysis. To satisfy these requirements, the compounds of the invention include: 1) An aryl ether group at position 6 of fluorescein which can be easily O-dealkylated by the enzyme, and 2) a fluorescein core for ease in fluorescence detection.


The enzyme reaction and assay method which describe the invention are illustrated in Reaction Scheme 1, below. Cytochrome P450 enzymes catalyze the dealkylation of compounds of Formula I to the intermediate of Formula II. Addition of base to the assay mixture hydrolyzes the ester intermediate (if present) to fluorescein which is quantitated spectrofluorometrically.

Reaction Scheme 1:



According to one aspect of the invention, compounds of Formula A are provided:

Formula A:




- (a) wherein R1 and R2 are independently selected from the group consisting of an hydrido and an aryl, provided that R1 and R2 are not both hydrido; and wherein the aryl contains an aryl ring carbon and/or an aryl ring nitrogen and the $(CH_2)_n$ or $(CH_2)_m$ is coupled via a covalent bond to the aryl ring carbon or the aryl ring nitrogen;
- (b) wherein n is 0, 1, 2, or 3;
- (c) wherein X1, X2, X3, X4, X5, and X6 are independently selected from the group consisting of an hydrido, a chloro, a fluoro, a bromo, and an iodo, provided that when R1 and R2 are phenyl, and n and m are 1, at least one of X1, X2, X3, X4, X5, and X6 is not hydrido;
- (d) wherein m is 0, 1, 2, or 3; and
- (e) wherein the compound is a cytochrome P450 substrate or a cytochrome P450 reaction product.

The above structure embraces three preferred embodiments, which are referred to herein as: (1) compounds of Formula I; (2) compounds of Formula II; and (3) compounds of Formula III. Compounds of Formula I and III are substrates for cytochrome P450 enzymes; compounds of Formula II are the products of a cytochrome P450 reaction in which a compound of Formula I is the substrate, i.e., the compounds of Formula II can be produced by the process of contacting a compound of Formula I with a cytochrome P450 under conditions whereby the cytochrome P450 enzyme catalyzes the conversion of the substrate to the reaction product.

According to certain embodiments, compounds of Formula I are provided:

Formula I:



(a) wherein R1 is aryl and R2 is independently selected from the group consisting of an hydrido and an aryl, provided that R1 and R2 are not both phenyl when n and m are 1; and wherein the aryl contains an aryl ring carbon and/or an aryl ring nitrogen and the $(CH_2)_n$ or $(CH_2)_m$ is coupled via a covalent bond to the aryl ring carbon or the aryl ring nitrogen;

5 (b) wherein n is 0, 1, 2, or 3;

(c) wherein X1, X2, X3, X4, X5, and X6 are independently selected from the group consisting of an hydrido, a chloro, a fluoro, a bromo, and an iodo;


(d) wherein m is 0, 1, 2, or 3; and

(e) wherein the compound is a cytochrome P450 substrate.

10 In the most preferred embodiments of the compound of Formula I, X2 and X5 are chloro and/or fluoro. In yet other embodiments, X2 and X5 may or may not be halogenated (particularly chloro and/or fluoro) and one or both of R1 and R2 are phenyl when one or both of n and m are 1. In still other preferred embodiments one or both of R1 and R2 are phenyl when one or both of n and m are 1, and the phenyl further includes (e.g., at the para position)
 15 an electron donating group, such as a halogen (especially chloro, fluoro), C (halogen)₃, or NH₃⁺.

According to yet other embodiments, compounds of Formula II are provided:

A3. The compound of claim A1, having **Formula II**:



20 (a) wherein R1 is hydrido and R2 is an aryl; and wherein the aryl contains an aryl ring carbon and/or an aryl ring nitrogen and the $(CH_2)_m$ is coupled via a covalent bond to the aryl ring carbon or the aryl ring nitrogen;

(b) wherein n is 0;

(c) wherein X1, X2, X3, X4, X5, and X6 are independently selected from the group


25 consisting of an hydrido, a chloro, a fluoro, a bromo, and an iodo;

(d) wherein m is 0, 1, 2, or 3; and

(e) wherein the compound is a cytochrome P450 reaction product.

For the most preferred embodiments of the compound of Formula II, X2 and X5 are chloro and/or fluoro. In yet other embodiments, X2 and X5 may or may not be halogenated (particularly chloro and/or fluoro), R2 is phenyl and m is 1. In still other preferred
 5 embodiments R2 is phenyl, m is 1, and the phenyl further includes (e.g., at the para position) an electron donating group, such as a halogen (especially chloro, fluoro), C (halogen)₃, or NH₃⁺.


A particularly preferred embodiment of the compound of Formula II is compound IIa:



Compound IIa

10 According to still other embodiments, compounds of Formula III are provided:

Formula III:



- (a) wherein R1 and R2 are each phenyl; and wherein the phenyl contains an aryl ring
 15 carbon and the (CH₂)_n or (CH₂)_m is coupled via a covalent bond to the aryl ring carbon;
- (b) wherein n is 1;
- (c) wherein X1, X2, X3, X4, X5, and X6 are independently selected from the group consisting of a chloro, a fluoro, a bromo, and an iodo;
- (d) wherein m is 1; and
- 20 (e) wherein the compound is a cytochrome P450 substrate.

For the most preferred embodiments of the compound of Formula III, X2 and X5 are chloro and/or fluoro. In yet other embodiments, X2 and X5 may or may not be halogenated


(particularly chloro and/or fluoro). In still other preferred embodiments R2 is phenyl and the phenyl further includes (e.g., at the para position) an electron donating group, such as a halogen (especially chloro, fluoro), C (halogen)₃, or NH₃⁺.

According to yet another aspect of the invention, a composition comprising a
5 compound of Formula I, a compound of Formula II, a compound of Formula III, and/or
Benzyloxyfluorescein benzyl ester ("DBF"; 97744-44-0; Benzoic acid, 2-[3-oxo-6-
(phenylmethoxy)-3H-xanthen-9-yl)-benzyl ester (CA index name), Hargreaves, J.S. and S.E.
Webber (1985) Can. J. Chem. 63, 1320-1327)) is provided. The compound is present in the
composition at a concentration greater than at least 50% by weight. The most preferred
10 compositions contain a concentration of the compound of Formula I, II, III, or DBF that is at
least 80%, more preferably at least 90%, and most preferably at least 95% by weight.
Preferably, the compositions are aqueous preparations which are free of contaminants that
would interfere with a cytochrome P450 enzyme reaction. Certain embodiments of the
preferred compositions are also substantially free of detectable reaction product, i.e., the
15 compositions of the invention do not contain levels of the cytochrome P450-catalyzed reaction
product. In these and other embodiments, the compositions optimally further contain a
cytochrome P450, e.g., an isolated enzyme or a microsome expressing a cDNA expressed
cytochrome P450.

The compositions may be contained in vials that are components of a kit for assaying
20 cytochrome P450 enzyme activity. The vials may contain preselected amounts of the
compositions to facilitate dissolution of the contents to achieve a preselected concentration of
the compound for performing a cytochrome P450 enzyme assay. Accordingly, in certain
embodiments of the invention, the compositions contain the appropriate buffers for performing
an enzyme reaction in which the compound of Formula I, the compound of Formula III, or
25 DBF serves as the substrate to form a fluorescent product.

According to yet another aspect of the invention, a method for assaying cytochrome
P450 enzyme activity is provided. The method involves contacting a cytochrome P450
enzyme with a compound of Formula I, a compound of Formula III, or DBF under conditions
whereby the cytochrome P450 enzyme catalyzes the conversion of the compound (substrate) to
30 a cytochrome P450 reaction product. Such conditions are generally known to those of ordinary
skill in the art and are illustrated in the Examples.

Reaction Scheme 1:



For ease of illustration, the above reaction is shown with respect to substrates which are compounds of Formula I as defined herein. It is to be understood that the reaction also can be performed wherein the substrate is a compound of Formula III as defined herein or DBF and that these compounds optionally are halogenated as shown in their respective formulas.

The method for assaying cytochrome P450 enzyme activity is used to detect activity of a cytochrome P450 that may be contained in biological fluid sample or solid sample (e.g., a biopsy sample from liver, brain or intestine) or that may be expressed in a cell-containing or cell-free system (e.g., a microsome containing cDNA-expressed cytochrome P450). In this manner, conditions associated with deficiencies or over expression of cytochrome P450 enzyme activity can be detected.

The cytochrome P450 assay may be performed in vivo or in vitro. For example, the compounds of the invention (e.g., the compounds of Formula I, the compounds of Formula III, or DBF) can be administered to an animal model for, e.g., locating and, optionally, quantifying, cytochrome P450 enzyme activity (e.g., by observing reaction products in biological fluid or tissue samples of the animal). More preferably, the method for assaying cytochrome P450 enzyme activity is used to detect activity of a cytochrome P450 that may be contained in biological fluid sample or solid sample (e.g., a biopsy sample from liver, brain or intestine) or that may be expressed in a cell-containing or cell-free system (e.g., a microsome containing

cDNA-expressed cytochrome P450). In this manner, conditions associated with deficiencies or over expression of cytochrome P450 enzyme activity can be detected. Thus, the cytochrome P450 enzyme may be contained in a sample that is a liver sample such as a crude homogenate, partially purified, or purified liver enzyme obtained from a biopsy, a cDNA-expressed

5 cytochrome P450, in hepatocytes, or in microsomes.

According to yet another aspect of the invention, a screening method for identifying agents which inhibit cytochrome P450 enzyme activity is provided. The method involves contacting a cytochrome P450 enzyme with a compound of Formula I, a compound of Formula III, or DBF in the presence of a putative cytochrome P450 enzyme inhibitor and identifying an agent which inhibits the cytochrome P450 enzyme activity as the cytochrome P450 enzyme inhibitor. In the preferred embodiments, the screening method is a high throughput screening assay. More preferably, the screening assay is performed in a multiwell (e.g., microtiter) plate or a container for containing a relatively small volume, e.g., the compound of Formula I, Formula III, or DBF is contacted with the putative cytochrome P450 enzyme inhibitor and the cytochrome P450 enzyme in a microtiter plate well or a small vial. In certain embodiments, the compounds of Formula I, Formula III, or DBF can be provided in the microtiter plate well or small vial. For example, the compounds can be distributed into one or more vials, which are then lyophilized or otherwise dried to provide a product having an enhanced shelf life. If the product is provided for use in a kit for measuring cytochrome P450 enzyme activity, the kit can further contain instructions for redissolving the compounds of Formula I, Formula III, or DBF and, optionally, an appropriate buffer (e.g., enzyme reaction buffer) for effecting the dissolution.

According to another aspect of the invention, a method for visualizing a cytochrome P450 enzyme is provided. The method involves contacting a cytochrome P450 enzyme-containing sample with a compound of Formula I, a compound of Formula III, or DBF and subjecting the cytochrome P450 enzyme and the compound to conditions whereby the cytochrome P450 enzyme catalyzes the conversion of the compound of Formula I, the compound of Formula III, or DBF to a fluorescent product. In the preferred embodiments, the method for visualizing a cytochrome P450 enzyme is performed on a tissue section sample, i.e., the cytochrome P450 enzyme-containing sample is a tissue section such as derived from a biopsy sample.

According to still another aspect of the invention, kits for detecting and/or measuring cytochrome P450 enzyme activity are provided. The kits contain a compound of Formula I, Formula III, and/or DBF and instructions for using the kits to measure cytochrome P450 enzyme activity. The kits may further contain instructions for calculating K_i and/or IC_{50} for a cytochrome P450 inhibitor. The preferred compounds of Formula I and compounds of Formula III are compounds which have a high specificity of binding for the enzyme and for which the enzyme exhibits a high rate of substrate turnover. These parameters typically are reflected in the K_m and V_{max} values for the enzyme-catalyzed conversion of the substrate (i.e., compound of the invention) to a fluorescent product. In general, a higher relative affinity of the cytochrome P450 enzyme for a first substrate compared to a second substrate is indicated by a lower K_m value for the first substrate compared to the second substrate. A higher catalytic turnover for a first substrate compared to a second substrate is indicated by a higher V_{max} for the first substrate. The preferred compounds of the invention have a K_m of greater than about 0.50 μM with a V_{max} greater than about 0.05 min^{-1} . In general, the compounds of the invention have a K_m from about 0.5 μM to about 5 μM and a V_{max} from about 0.10 or from about 0.20 to about 50 min^{-1} , with a preferred range for K_m being greater than about 0.50 μM and a preferred range for V_{max} being from about 0.20 min^{-1} to about 50 min^{-1} ; with V_{max} being more preferably from about 2.0 or from about 20.0 to about 50 min^{-1} .

According to yet another aspect of the invention, novel cytochrome P450 fluorescent products are provided. The novel products are compounds of Formula A wherein R1 is hydrido and n is 0 (see Formula II). In general, these novel products are produced as a reaction product of the cytochrome P450-catalyzed reaction of a substrate that is a compound of Formula I, a compound of Formula III, or DBF. In general, these compounds have structures that differ from those of the compounds of Formula I, Formula III, and DBF in having a hydroxy group at position 6 of the fluorescein.

These and other aspects of the invention as well as various advantages and utilities will be more apparent with reference to the detailed description of the preferred embodiments. All patents, patent publications and references identified in this document are incorporated in their entirety herein by reference.

Brief Description of the Drawings

The drawings are illustrative only and are not required for enablement of the claimed invention.

Fig. 1 shows selectivity for Compound 1a dealkylation by a panel of Human P450 enzymes and Effect of Delay in Read Time following addition of 2N Stop Solution on Signal to Noise Ratio.

Fig. 2 shows selectivity for Compound 1b dealkylation by a panel of Human P450 enzymes and Effect of Delay in Read Time following addition of 2N Stop Solution on Signal to Noise Ratio.

Fig. 3A shows selectivity for CEC* dealkylation in a panel of human P450 enzymes.
Fig. 3B shows selectivity for MFC* dealkylation in a panel of human P450 enzymes.
Fig. 3C shows selectivity of 7-BQ* dealkylation in a panel of human P450 enzymes .
Fig. 3D shows selectivity for BFC* dealkylation in a panel of human P450 enzymes.
Fig. 3E shows selectivity for AMMC* dealkylation in a panel of human P450 enzymes.
Fig. 3F shows selectivity for dealkylation of Benzyloxyresorufin in a panel of human P450 enzymes.

Fig. 4A shows selectivity for 5-(and-6)-chloromethylfluorescein diethyl ether (CMFDE) dealkylation by a panel of Human P450 Enzymes.

Fig. 4B shows selectivity for 3-O-methylfluorescein (MF) by a panel of Human P450 Enzymes.

Detailed Description of the Invention

I. Definitions

Throughout this document cytochrome P450 is used in reference to the enzyme which catalyzes the conversion of a compound of the invention to a fluorescent product. It is to be understood that any member of the cytochrome P450 family can be used in any of the enzyme reactions discussed herein and that CYP2C8, CYP2C9, CYP2C19, and CYP3A4 represent a particularly preferred embodiment of the invention.

Molecular terms, when used in this application, have their common meaning unless otherwise specified. The term hydrido denotes a single hydrogen atom.

Aryl groups can contain from 0-4 hetero atoms selected from oxygen, nitrogen and sulfur in a single or fused carbocyclic or heterocyclic ring system, having from 5-15 ring

members. One or more hydrogen atoms may also be replaced by a substituent group selected from an acyl, an amino, a carboalkoxy, a carboxy, a carboxyamido, a cyano, a halo, a hydroxy, a nitro, a thio, an alkyl, an aryl, a cycloalkyl, an alkoxy, an aryloxy, a sulfoxy, and a guanido group. When the aryl group is coupled to a linker such as $(CH_2)_n$ or $(CH_2)_m$, a carbon or a
5 nitrogen of the aryl ring is covalently coupled to the linker.

A preferred class of aryl groups are unsubstituted phenyl groups and phenyl groups in which one or more hydrogen have been replaced with an alkyl, alkoxy, aryloxy, or halo group. Exemplary aryl groups include phenyl, phenyl naphthyl, biphenyl, terphenyl, pyridinyl, and various other phenyl derivatives.

10

II. Description

The invention provides compounds of Formula A and methods for manufacturing and/or using the compounds of Formula I, II, III, and DBF for assaying the activity of cytochromes P450. The compounds are particularly useful for measuring the potential
15 inhibition of cytochromes P450, preferably in a high throughput screening assay. For example, the invention provides a method for assaying cytochromes P450 which involves contacting a cytochrome P450 enzyme with a compound of Formula I, a compound of Formula III, or DBF under the conditions in which the cytochrome P450 enzyme interacts with the compound and catalyzes dealkylation at the 6 position of the compound to form a 6-hydroxy-fluorescein ester
20 product. Such conditions are known to those skilled in the art (see also, e.g., the Examples for conditions). This method can be performed using in vivo or in vitro sources of cytochrome P450 enzyme. In a further aspect, the invention provides a method for assessing the potential cytochrome P450 inhibition of a test chemical, preferably in a high throughput screening assay.

The structures of the compounds of the invention, particularly the structures of Formula
25 A, Formula I, Formula II, and Formula III, are provided in the Summary and in the Claims.


General Synthetic Procedures

For ease of illustration, the reaction illustrated below are shown with respect to substrates which are compounds of Formula I. It is to be understood that the reaction also can
30 be performed wherein the substrate is a compound of Formula III or DBF and that these compounds optionally are halogenated as shown in their respective formulas.

General Procedure 1

Compounds of Formula I and Formula III are synthesized based on the method of Hargreaves and Webber (Can. J. Chem. 63, 1320-1327 (1985)). The commercially available reagent, fluorescein (compound 1), is converted to a compound of Formula I or a compound of

5 Formula III by treatment with compound 2 and reagent A in an appropriate solvent such as tetrahydrofuran, acetone, dimethyl sulfoxide, acetonitrile, or dimethylformamide at temperatures ranging from 0°C to 100°C.


Reaction 1

- 10 wherein R¹, R², n, and m are defined as above; wherein A is a base such as potassium carbonate or lithium hydroxide; and X is a leaving group such as bromide, chloride, iodide, or tosylate.

General Procedure 2

A compound of Formula I, is converted to another compound of Formula I by treatment

15 with reagent B in an appropriate organic solvent such as tetrahydrofuran, acetone, dimethylsulfoxide, acetonitrile, or dimethylformamide mixed with water at temperatures ranging from 0°C to 100°C. A preferred compound for performing this reaction is benzyloxyfluorescein benzyl ester (compound IIa, also referred to herein as "DBF") to yield a preferred compound of Formula I, Compound Ib (MBF).

Reaction 2

wherein R1 is defined as above; wherein B is a base such as sodium hydroxide or lithium hydroxide.

Examples

5 The following examples are detailed descriptions of the methods of making and using the compounds of Formula A. These detailed preparations fall within the scope of, and serve to exemplify, the above described General Procedures which form part of the invention. These Examples are presented for illustrative purposes only and are not intended to limit the scope of the invention.

10 **Example 1: Preparation of O-Benzylfluorescein Benzyl Ester (DBF). (Compound Ia)**



Fluorescein (40 g, 120.5 mmol) was dissolved in dimethylformamide (DMF, 500 mL) in a 2 L flask fitted with a mechanical stirrer then potassium carbonate (72 g, 521.7 mmol) was added in as a single portion. The reaction was heated to 80°C then benzyl chloride (42 mL, 363.5 mmol) was added dropwise. After 24 h at 80°C the reaction was cooled down to room
15 temperature. Acetone (400 mL) was added and the product was allowed to precipitate overnight. The solid orange product was filtered, washed with acetone and dried in a vacuum oven at 50°C. The O-benzylfluorescein benzyl ester was obtained as an orange solid (36.4 g, 59%): mp 192-193°C (Lit. 193-194°C); ¹H NMR (DMSO-d₆): 4.97 (dd, 2H, J = 12.2 Hz, J = 15.3 Hz), 5.28 (s, 2H), 6.13 (d, 1H, J = 1.9 Hz), 6.36 (dd, 1H, J = 1.9 Hz, J = 9.9 Hz), 6.77 (d,
20 1H, J = 9.9 Hz), 6.82 6.77 (d, 1H, J = 9.0 Hz), 6.90-6.98 (m, 3H), 7.14-7.27 (m, 4H), 7.33-7.52 (m, 6H), 7.78 (dt, 1H, J = 1.65 Hz, J = 7.4 Hz), 7.85 (dt, 1H, J = 1.65 Hz, J = 7.4 Hz), 8.23 (dd, 1H, J = 1.35 Hz, J = 7.7 Hz); ¹³C NMR (DMSO-d₆): 67.4, 70.7, 101.9, 105.2, 114.7, 115.0, 117.3, 128.3, 128.5, 128.7, 128.8, 128.9, 129.4, 130.0, 130.2, 130.6, 130.8, 131.2, 131.4, 133.8, 134.1, 135.3, 136.6, 150.1, 153.9, 158.7, 163.3, 165.5, 184.4; IR (nujol) cm⁻¹ 1728, 1643,
25 1595, 1517, 1462, 1377, 1261, 1211, 1081, 855. C₃₄H₂₄O₅: Calc.(C, 79.67; H, 4.72) Found (C, 79.74; H, 4.79).

Example 2: Preparation of Monobenzyl fluorescein (MBF) Compound Ib.

O-Benzylfluorescein benzyl ester, compound Ia (20.0 g, 39 mmol), was dissolved in THF/H₂O 10/1 (400 mL) and lithium hydroxide monohydrate (6.0 g, 158 mmol) was added. The reaction was refluxed for two hours then cooled down to room temperature. The lithium salt was removed by filtration, and the filtrate was diluted in water and extracted with methylene chloride. The organic phase was dried over magnesium sulfate and evaporated under reduced pressure followed by drying in a vacuum oven at 60°C. The monobenzyl fluorescein was obtained as a yellow solid (11.9 g, 70%): mp 113-115°C; ¹H NMR (DMSO-d₆): 5.17 (s, 2H), 6.58 (s, 2H), 6.64-6.71 (m, 2H), 6.68 (dd, 1H, J = 2.2 Hz, J = 8.8 Hz), 7.02 (d, 1H, J = 2.2 Hz), 7.26-7.47 (m, 6H), 7.72 (t, 1H, J = 7.2 Hz), 7.79 (t, 1H, J = 7.4 Hz), 8.01 (d, 1H, J = 7.7 Hz), 10.18 (s, 1H); ¹³C NMR (DMSO-d₆): 70.2, 83.2, 102.2, 102.8, 110.0, 111.8, 113.1, 113.4, 124.6, 125.2, 126.6, 128.3, 128.6, 129.0, 129.6, 129.7, 130.7, 136.2, 137.1, 152.3, 153.0, 160.1, 160.6, 169.2; IR (nujol) cm⁻¹ 1727, 1609, 1501, 1462, 1378, 1286, 1249, 1172, 1108. C₂₇H₁₈O₅; Calc.(C, 76.28; H, 4.34) Found (C, 76.22; H, 4.32).

TABLE I

#	Structure	General Procedure
Ia		1
Ib		2

**Compound Ia****Compound Ib**

B. Biological Evaluation of Compound I.

The evaluation of compounds of Formula I for usefulness as cytochrome P450 substrates includes: (1) The specificity of the compounds of Formula I as substrates for individual cytochromes P450 was examined using cDNA-expressed cytochromes P450; (2) 5 Enzyme kinetics for the compounds of Formula I were performed using cDNA-expressed cytochromes P450; and (3) The IC_{50} values for known inhibitors of cytochromes P450, measured in a high throughput screening assay, of compounds of Formula I, were correlated with those values from known substrates useful for this application.

(1) Substrate Specificity

10 The specificity of several cytochrome P450 enzymes for the dealkylation of Compound Ia (Figure 1) and Compound Ib (Figure 2) was examined. Compounds of Formula I are substrates for multiple isoforms within the three major families of drug metabolizing P450 enzymes (as demonstrated by a high signal to noise ratio). While it is well known that fluorescein fluorescence is best detected at high pH (>9), a novel aspect of this assay, which 15 contrasts with previously described fluorescence-based assays, is that the signal to noise ratio was dramatically increased when the microtiter plate readings were taken 2 hours after the reaction was stopped by the addition of 2N sodium hydroxide solution (2N NaOH).

CYP1A1 has the highest activity with compound Ia. CYP2C8 also exhibits substantial enzyme activity for Compound Ia. This is the first fluorescent substrate ever reported for 20 CYP2C8. CYP2C9*1, CYP2C19, and members of the CYP3A family also catalyzed the dealkylation of compound Ia. Compound Ia is a more sensitive substrate (higher signal to noise ratio) for CYP2C9 and CYP2C19 compared to the previously described alkylcoumarin fluorescent substrates. Compound Ib as a substrate for the various P450s gave substantially less signal versus Compound Ia and exhibited roughly a similar enzyme specificity profile. For 25 the fluorometric assay of cytochromes P450 CYP1A1, CYP2C8, CYP2C9, CYP2C19, and CYP3A, compound Ia is clearly superior to compound Ib.

Example Compounds Ia and Ib. Assays were conducted in 96 well microtiter plates (Corning COSTAR, cat. no. 3915). The substrate, compound Ia, was prepared in acetonitrile. After cofactors addition, the plates were prewarmed to 37°C. Incubations were initiated by the 30 addition of prewarmed enzyme and substrate. The enzymes were commercially available, baculovirus/insect cell expressed human P450s (SUPERSOMES®, GENTEST Corporation).

The amount of enzyme added per well was 2 pmole. The final cofactor concentrations were 1.3 mM NADP, 3.3 mM glucose-6-phosphate and 0.4 U/mL glucose-6-phosphate dehydrogenase. Final incubation volume was 0.2 ml. Incubations were carried out for 30 minutes and stopped by the addition of 0.075 mL of 2 N sodium hydroxide solution.

5 Fluorescence per well was measured using a BMG FLUOstar fluorescence plate scanner equipped with an IBM-compatible computer. The metabolite was measured using an excitation wavelength of 485 nm and emission wavelength of 530 nm. Data was exported and analyzed using an Excel spreadsheet.

(2) Enzyme Kinetics

10 The evaluation of compounds of Formula I were performed by measuring the kinetics of turnover using cDNA-expressed cytochromes P450 CYP3A4 and CYP2C8. V_{max} and K_m values are important for the optimization of the assay conditions and for setting the parameters for inhibition experiments. Fluorometric assays for the turnover of compounds of Formula I by cytochromes P450 were performed based on a modification of the method by Crespi et al. *Anal*
15 *Biochem.* **248**, 188-190, (1997).

Comparisons of the CYP3A4 enzyme kinetics for compound Ia and the fluorescent substrate, 7-benzyloxyresorufin (BzRes), show that CYP3A4 has a higher affinity for compound Ia (lower K_m) and a higher catalytic turnover for compound Ia (higher V_{max}) (See Table II). Additionally, compound Ia is a more robust substrate in terms of solubility and
20 product detection properties. Based on these results, compound Ia is a better fluorescent substrate for CYP3A4 than the previously described BzRes.


TABLE II

Substrate	Enzymes	K_m (uM)	V_{max}
Ia	CYP3A4	0.87 ± 0.12	22 ± 0.98
BzRes	CYP3A4	30 ± 7	2.3 ± 0.2

For CYP2C8, the K_m was found to be 1.4 uM and the V_{max} was determined to be 0.22
25 min^{-1} . These parameters are reasonable allowing this assay to be utilized for high throughput fluorescent assays using microtiter plates. As stated earlier, this is the first fluorescent substrate reported for this important human cytochrome P450 enzyme.


CYP2C8 is the human cytochrome P450 primarily responsible for the metabolism of

the anti-cancer drug paclitaxel (TAXOL™, Rahman, et al., 1994. *Cancer Research*. **54**, 5543-5546). The structure of paclitaxel and the site of hydroxylation by CYP2C8 is shown in the figure below. Notice that this large complicated molecule does not resemble any of the fluorescent substrates described herein, including DBF.



5

In addition, the following fluorescent molecules have been tested as a substrate probe for CYP2C8 and the assay results obtained for these substrates (Figure 3) further illustrate the unexpected and surprising specificity of cytochrome P450 CYP2C8 for DBF:



10

Data for these specificity experiments is shown in Figure 3 (panels 3A, 3B, 3C, 3D, 3E, and 3F).

Assays were conducted in 96-well microtiter plates. Microsomes were obtained from GENTEST Corp from metabolically competent human B-lymphoblastoid cell lines that stably express rat CYP2E1 or rat CYP2A1 or from baculovirus-infected insect cells

(SUPERSOMES®, all other enzymes). Substrates and metabolites were obtained from

- 5 GENTEST Corporation except fluorescein which was obtained from Sigma-Aldrich. All other chemicals (reagent grade) were obtained from Sigma-Aldrich. The substrates and metabolites examined and the wavelength used for analysis are shown in **Table III**. Potassium phosphate buffer (0.1 mL) containing an NADPH-regenerating system was added to each well. The plate was then warmed to 37 °C and the reaction initiated by the addition of pre-warmed
- 10 enzyme/substrate (E/S) mix. The E/S mix contained buffer, cDNA-expressed P450 (2.5-87 pmol/mL final), control protein from mock-transfected insect cells (in order to standardize protein to approximately 0.25 mg/mL final) and substrate (1.0 µM-50 µM final). Reactions were terminated after 20-45 min by addition of 75 µl 80:20 acetonitrile:0.5 M Tris base or for Compound Ia only, 2 N NaOH. The final concentration for all enzymes was 50 nM except for
- 15 CYP2A1 (20 nM) and for CYP2E1 (87 nM). Fluorescence signal was measured using a FLUOstar model 403 fluorescence plate reader (BMG LabTechnologies, Inc., Durham, NC).

TABLE III

Fluorometric substrates and their metabolite excitation and emission wavelengths				
Substrate	Substrate Abbreviation	Metabolite	Ex (nm)	Em (nm)
3-[2-(N,N-Diethyl-N-methylamino) ethyl]-7-methoxy-4-methylcoumarin	AMMC	3-[2-(N,N-Diethyl-N-methylamino) ethyl]-7-hydroxy-4-methylcoumarin	390	460
3-[2-(N,N-Diethyl-N-methylamino) ethyl]-7-methoxy-4-trifluoromethylcoumarin	MeAMFC	3-[2-(N,N-Diethyl-N-methylamino) ethyl]-7-hydroxy-4-trifluoromethylcoumarin	410	538
7-Methoxy-3-cyanocoumarin	CMC	7-hydroxy-3-cyanocoumarin	410	460
7-Ethoxy-3-cyanocoumarin	CEC	7-hydroxy-3-cyanocoumarin	410	460
7-Methoxy-4-trifluoromethylcoumarin	MFC	7-hydroxy-4-trifluoromethylcoumarin	410	538
7-Benzyloxy-4-trifluoromethylcoumarin	BFC	7-hydroxy-4-trifluoromethylcoumarin	410	538
Compound Ia	Ia	Fluorescein	485	538
7-Benzyloxyquinoline	BQ	7-hydroxyquinoline	410	538
Resorufin benzyl ether	BzRes	Resorufin	530	590

Example Compound Ia. Assays were conducted in 96 well microtiter plates (Corning COSTAR, cat. no. 3915). The substrate, compound Ia, was prepared in acetonitrile. After cofactors addition, the plates were prewarmed to 37°C. Incubations were initiated by the addition of prewarmed enzyme and substrate. The enzymes were commercially available, baculovirus/insect cell expressed human CYP3A4 or CYP2C8 (SUPERSOMES®, GENTEST Corporation). The amount of enzyme added per well was 0.2 pmoles (CYP3A4) or 2 pmole (CYP2C8). The final cofactor concentrations were 1.3 mM NADP, 3.3 mM glucose-6-phosphate and 0.4 U/mL glucose-6-phosphate dehydrogenase. Final incubation volume was 0.2 ml. Incubations were carried out for 10 minutes (CYP3A4) or 30 min (CYP2C8) and stopped by the addition of 0.075 mL of 2 N sodium hydroxide solution. Fluorescence per well was measured using a BMG FLUOstar fluorescence plate scanner equipped with an IBM-compatible computer. The metabolite was measured using an excitation wavelength of 485 nm and emission wavelength of 530 nm. Data was exported and analyzed using an Excel spreadsheet. The activity was quantitated by comparing to a standard curve of fluorescein. Apparent Km and Vmax values were calculated by non-linear regression using SigmaPlot software.

(3) Example I. High Throughput Cytochrome P450 Inhibition Screen – CYP3A4

Measuring the cytochrome P450 inhibition potential of compounds of Formula I was based on a modification of the published method by Crespi et al. Anal Biochem. **248**, 188-190, (1997). With CYP3A4, a series of test compounds was examined in the modified system, using BzRes, 7-benzyloxy quinoline (BQ), benzyloxy trifluoromethyl coumarin (BFC) or compound Ia as a substrate (Table IV). In general, inhibitors of P450 enzymes exhibit similar IC₅₀ values independent of substrate. However, this is not the case with CYP3A4. A several fold range in IC₅₀ values was found and these results are consistent with the unusual kinetics associated with this enzyme (KE Thummel and GR Wilkinson, Ann. Rev. Pharmacol. Toxicol. **38**:389-430 (1998)). One explanation is that the CYP3A4 enzyme is capable of accommodating 2 or more compounds simultaneously, one of which may activate, or inhibit metabolism of the other. The use of multiple substrates (with diverse chemical properties, i.e. size, shape, etc.) to test for inhibition of CYP3A4 has been recommended. Thus, compound Ia, represents a valuable tool for exploring the potential for a drug to interact with other, co-administered drugs and

CYP3A4. Moreover, the utility of Compound Ia as a CYP3A4 substrate that utilizes only small amounts of enzyme to perform the assay was also demonstrated.

The utility of compound Ia for measuring the inhibition of CYP2C8 is demonstrated in Table V. A series of 8 test compounds was examined. Values obtained are consistent with those expected. For example, sulfaphenazole exhibited an elevated IC_{50} , compared to the IC_{50} obtained with CYP2C9 (~0.22 μ M), another member of the CYP2C family. Additionally, the prototypical substrate for the enzyme, paclitaxel, inhibited the turnover of compound Ia which is consistent with the conclusion that these two molecules are competing substrates for CYP2C8.

Table IV. Absolute and relative Mean IC_{50} Values for CYP3A4 with Compound Ia substrate (μ M)

Test Compound	Absolute Mean IC_{50} Values						Relative Mean Values			
	BzRes	BQ	BFC	Ia	Overall	Range	BzRes	BQ	BFC	Ia
Itraconazole	0.176	0.350	0.034	0.038	0.176	10	1.000	1.987	0.195	0.216
Ketoconazole	0.034	0.072	0.005	0.002	0.028	46	1.198	2.563	0.183	0.056
Miconazole	0.175	0.613	0.057	0.058	0.226	11	0.775	2.718	0.252	0.256
Erythromycin	1.797	54.443	4.496	25.344	21.520	30	0.084	2.530	0.209	1.178
Cisapride	0.107	0.022	0.022	0.922	0.268	41	0.398	0.083	0.083	3.436
Mibefradil	0.021	0.066	0.005	0.054	0.037	12	0.561	1.807	0.146	1.486
Midazolam	59.930	3.560	1.250	1.687	16.607	48	3.609	0.214	0.075	0.102
Clotrimazole	0.019	0.013	0.002	0.005	0.010	7.9	1.943	1.285	0.247	0.525
Nifedipine	68.494	16.782	8.383	4.132	24.448	17	2.802	0.686	0.343	0.169
Cyclosporin	1.200	146.885	3.122	3.307	38.629	122	0.031	3.803	0.081	0.086
Verapamil	0.398	8.401	0.357	4.190	3.336	21	0.119	2.518	0.107	1.256
Terfenadine	Activation	6.307	1.001	2.149	3.152	6.3	-	2.001	0.318	0.682
a-Naphthoflavone	Activation	Activation	Activation	146.148	-	-	-	-	-	-
Tamoxifen	Activation	>1000	>1000	>1000	-	-	-	-	-	-
Carbamazepine	Activation	478.634	174.969	89.943	247.849	5.3	-	1.931	0.706	0.363
Ethinylestradiol	6.401	3.752	1.425	3.319	3.724	4.5	1.719	1.008	0.383	0.891
Testosterone	Activation	>1000	Activation	32.108	-	-	-	-	-	-
TAO	0.292	6.105	0.412	1.925	2.184	21	0.134	2.796	0.189	0.882
Progesterone	Activation	> 200	Activation	11.501	-	-	-	-	-	-
Quercetin	> 50	> 50	> 50	> 50	-	-	-	-	-	-
					Mean	27	1.105	1.862	0.234	0.772

Table V. IC_{50} Values with Compound Ia as a CYP2C8 Substrate (μ M)



Test Compound	IC_{50}
Quercetin	2.35
Paclitaxel	13.5
Sulfaphenazole	182
Carbamazepine	> 50
Phenytoin	> 50
Ketoconazole	6.65
Cyclosporin	> 200
Diclofenac	> 50

Example Compound Ia. Assays were conducted in 96 well microtiter plates. The substrates, compounds of Formula I, were prepared in acetonitrile. The substrate stock concentrations were twice the final concentration (final concentration chosen to be approximately the apparent K_m , for example 1-2 μ M for compound I). The 12 wells in a row were used for one test. Wells 1 to 8 contained serial 1:3 dilutions of the inhibitors. Wells 9 and 10 contained no inhibitor and rows 11 and 12 were blanks for background fluorescence (stop solution added before the enzyme). After substrate and inhibitor addition, the plates were prewarmed to 37°C. Incubations were initiated by the addition of prewarmed enzyme and cofactors. The enzymes were commercially available, baculovirus/insect cell expressed human CYP3A4 (SUPERSOMES®, Cat. no. P202, GENTEST Corporation) or CYP2C8 (SUPERSOMES®, Cat. no. P252, GENTEST Corporation). The amount of enzyme added per well was 0.2 pmoles (CYP3A4) or 4 pmoles (CYP2C8). The final cofactor concentrations were 1.3 mM NADP, 3.3 mM glucose-6-phosphate and 0.4 U/mL glucose-6-phosphate dehydrogenase. Final incubation volume was 0.2 ml. Incubations were carried out for 10 minutes (CYP3A4) or 30 min (CYP2C8) and stopped by the addition of 2 N sodium hydroxide solution. Fluorescence per well was measured using a BMG FLUOstar fluorescence plate reader controlled with an IBM-compatible computer. The metabolite was measured using an excitation wavelength of 485 nm and emission wavelength of 530 nm. Data was exported and analyzed using an Excel spreadsheet. The IC_{50} values were calculated by linear interpolation.

Example 3: Specificity of Other Alkyl Fluorescein Compounds

The specificity of several cytochrome P450 enzymes for the dealkylation of two different commercially available alkyl ether fluoresceins was examined. 5-(and 6)-Chloromethylfluorescein diethyl ether (CMFDE, a mixture of isomers) is very similar to the diethoxyfluorescein studied by A. G. Miller (A. G. Miller, Anal. Biochem. 133, 46-57 (1983)). CMFDE must be dealkylated twice in order to form the fluorescent fluorescein product -- 5(and 6)-carboxyfluorescein. A. G. Miller reported that diethyl ethers of fluorescein are poor P450 substrates in mouse liver microsomes because they must be metabolized twice by the P450 enzyme in order to generate the fluorescent product. 3-O-Methylfluorescein (MF) is a simple, single methyl ether of fluorescein. The use of MF as a P450 substrate has never been reported.

The specificity for dealkylation of CMFDE by human cytochromes P450 is summarized in Figure 4A. CMFDE is not a good substrate (based on turnover number) for any of the human P450 enzymes. CMFDE was a substrate for multiple P450 isoforms of drug metabolizing P450 enzymes showing little specificity except for CYP1A1. This metabolism profile is very different from that of other alkylfluoresceins (Compound Ia, Ib, and MF). The specificity for dealkylation of MF by human cytochromes P450 is summarized in Figure 4B. MF is a good P450 substrate (based on turnover number) for CYP1A1 and CYP2C19. Since CYP1A1 is rare in human liver, this substrate may be very useful as a CYP2C19 specific probe in human liver tissue. Both of these examples illustrate the difficulty in predicting human P450 specificity for the dealkylation of alkylfluoresceins. The specificity profiles are very different from those of Compounds Ia and Ib.



Examples CMFDE and MF. Assays were conducted in 96 well microtiter plates (Corning COSTAR, cat. no. 3915). The substrate, CMFDE or MF, was prepared in acetonitrile. After cofactors addition, the plates were prewarmed to 37°C. Incubations were initiated by the addition of prewarmed enzyme and substrate. The enzymes were commercially available, baculovirus/insect cell expressed human P450s (SUPERSOMES®, GENTEST Corporation). The amount of enzyme added per well was 2 pmole. The final substrate concentration was 1 μ M, final cofactor concentrations were 1.3 mM NADP, 3.3 mM glucose-6-phosphate and 0.4 U/mL glucose-6-phosphate dehydrogenase. Final incubation volume was 0.2 mL. Incubations were carried out for 30 minutes and stopped by the addition of 0.075 mL of 2 N sodium hydroxide solution. Fluorescence per well was measured using a BMG FLUOstar fluorescence plate scanner equipped with an IBM-compatible computer. The metabolite was measured using an excitation wavelength of 485 nm and emission wavelength of 530 nm and compared

to a standard curve of metabolite standard (5(and 6)-carboxyfluorescein for CMFDE and fluorescein for MF). Data was exported and analyzed using an Excel spreadsheet.

Example 4: Substrate Specificity with Rat P450s

Rats are a common *in vivo* animal model for human xenobiotic metabolism. There is much current interest in the species differences between rats and humans, especially *in vitro* models that would explain species differences or validate the rat as a good model for human xenobiotic metabolism. Few fluorescent assays have been developed for rat P450 enzymes. Most rat P450 assays are based on tedious HPLC assays. A quick and simple fluorescent plate-based assay would find utility in studying species differences in xenobiotic metabolism.

The specificity of several rat cytochrome P450 enzymes for the dealkylation of Compound Ia (Figure 1) and other fluorescent cytochrome P450 substrates was examined. The data (turnover number) is summarized in Table V. Compound Ia was a substrate for multiple isoforms within the three major families of drug metabolizing P450 enzymes (as demonstrated by a high signal to noise ratio). The catalytic profiles for each substrate, including Compound Ia, are distinct. For those rat P450s that turnover Compound Ia, Compound Ia is the most sensitive assay reagent based on its low limit of detection. This characteristic is likely due to the superior wavelength and quantum yield properties of the fluorescein product compared to the other coumarins and resorufin products. This data shows that Compound Ia is a promising fast and inexpensive substitute for many HPLC-based rat P450 assays.

Example Compound Ia. Assays were conducted in 96 well microtiter plates (Corning COSTAR, cat. no. 3915). The substrate, compound Ia, was prepared in acetonitrile. After cofactors addition, the plates were prewarmed to 37°C. Incubations were initiated by the addition of prewarmed enzyme and substrate. The enzymes were commercially available, baculovirus/insect cell expressed rat P450s (SUPERSOMES®, GENTEST Corporation). The amount of enzyme added per well was 2 pmole. The final cofactor concentrations were 1.3 mM NADP, 3.3 mM glucose-6-phosphate and 0.4 U/mL glucose-6-phosphate dehydrogenase. Final incubation volume was 0.2 mL. Incubations were carried out for 30 minutes and stopped by the addition of 0.075 mL of 2 N sodium hydroxide solution. Fluorescence per well was measured using a BMG FLUOstar fluorescence plate scanner equipped with an IBM-compatible computer. The metabolite was measured using an excitation wavelength of 485 nm and emission wavelength of 530 nm. Data was exported and analyzed using an Excel

spreadsheet.

Example Other Fluorescent P450 Substrates. The substrate, buffer and protein concentrations and incubation times (see Table VI) were chosen based primarily on those conditions determined to be optimal for cytochrome P450 inhibition experiments for the human enzyme (e.g. CYP3A4 with BFC). This refers to a substrate concentration at or near the apparent K_m and conditions of where metabolite production was linear with time and protein. Assay conditions were not necessarily optimal with all enzyme/substrate pairs, however, in most cases, substrate utilization was below 15%. A notable exception was CYP2D2/AMMC where substrate utilization was 45%. Therefore, the velocities reported were not necessarily measured under initial rate conditions.

TABLE VI

Turnover Numbers for Rat CYP Enzymes with Fluorometric Substrates ^a							
Rat CYP Isoform	Ia	AMM C	Substrate				
			CEC	MFC	BFC	BQ	BzRes
CYP1A1	0.211	- ^b	78	1.03	0.34	5.7	1.35
CYP1A2	0.041	-	1.38	1.32	-	2.3	0.62
CYP2A1 ^c	-	-	-	-	-	-	-
CYP2A2	0.030	-	0.04	0.08	0.26	2.8	-
CYP2B1	-	-	0.45	2.82	3.70	2.5	0.82
CYP2C6	0.021	-	2.14	11.5	2.62	1.9	0.02
CYP2C11	0.065	-	0.55	5.57	0.34	-	0.05
CYP2C12	-	-	-	0.24	-	1.5	-
CYP2C13	0.003	-	-	0.21	0.74	-	0.01
CYP2D1	0.159	-	-	0.09	0.25	5.3	0.02
CYP2D2	0.194	5.9	-	0.08	0.12	4.4	-
CYP2E1 ^c	-	-	0.10	2.11	-	-	-
CYP3A1	0.027	-	0.08	-	1.33	8.4	0.06
CYP3A2	0.015	-	0.15	-	1.04	9.8	0.09
Substrate concentration (uM)	2	20	40	50	50	40	50
Incubation time (min)	30	30	30	30	30	30	30
Enzyme concentration (nM) ^d	50	50	50 ^e	50	50	50	50

^a – Values, expressed as pmol product/min/pmol enzyme, represent the mean of duplicate

determinations. Bold values indicate the highest catalytic activity within the panel.

^b – Dashes indicate activity was below the limit of detection which was considered as that corresponding to a signal to background ratio of approximately 1.5. These values and their substrates were 0.05 min⁻¹, AMMC; 0.05 min⁻¹, BFC; 0.008 min⁻¹, CEC; 0.003 min⁻¹, Ia; 0.05 min⁻¹, MFC; 1.0 min⁻¹, BQ; 0.008 min⁻¹, BzRes.

^c – Microsomes from human lymphoblastoid cells engineered to express the rat enzyme.

^d – Concentration for all enzymes was 50 nM except for CYP2A1 (20 nM) and for CYP2E1 (87 nM).

^e – Concentration for all enzymes was 50 nM except for CYP1A1, which was 2.5 nM.


The preceding is merely a detailed description of certain preferred embodiments. It therefore should be apparent to those skilled in the art that various modifications and equivalents can be made without departing from the spirit or scope of the invention.

All references disclosed herein are incorporated by reference in their entirety.

5 What is claimed is:

CLAIMS

1. A compound of the formula:



(a) wherein R1 and R2 are independently selected from the group consisting of an
5 hydrido and an aryl, provided that R1 and R2 are not both hydrido; and wherein the aryl
contains an aryl ring carbon and/or an aryl ring nitrogen and the $(CH_2)_n$ or $(CH_2)_m$ is coupled
via a covalent bond to the aryl ring carbon or the aryl ring nitrogen;

(b) wherein n is 0, 1, 2, or 3;


(c) wherein X1, X2, X3, X4, X5, and X6 are independently selected from the group
10 consisting of an hydrido, a chloro, a fluoro, a bromo, and an iodo, provided that when R1 and
R2 are phenyl, and n and m are 1, at least one of X1, X2, X3, X4, X5, and X6 is not hydrido;

(d) wherein m is 0, 1, 2, or 3; and

(e) wherein the compound is a cytochrome P450 substrate or a cytochrome P450
reaction product.

15

2. The compound of claim 1, having **Formula I**:



(a) wherein R1 is aryl and R2 is independently selected from the group consisting
of an hydrido and an aryl, provided that R1 and R2 are not both phenyl when n and m are 1;
and wherein the aryl contains an aryl ring carbon and/or an aryl ring nitrogen and the $(CH_2)_n$ or
20 $(CH_2)_m$ is coupled via a covalent bond to the aryl ring carbon or the aryl ring nitrogen;

(b) wherein n is 0, 1, 2, or 3;


(c) wherein X1, X2, X3, X4, X5, and X6 are independently selected from the group consisting of an hydrido, a chloro, a fluoro, a bromo, and an iodo;

(d) wherein m is 0, 1, 2, or 3; and

(e) wherein the compound is a cytochrome P450 substrate.

5

3. The compound of claim 1, having **Formula II**:



(a) wherein R1 is hydrido and R2 is an aryl; and wherein the aryl contains an aryl ring carbon and/or an aryl ring nitrogen and the $(CH_2)_m$ is coupled via a covalent bond to the aryl ring carbon or the aryl ring nitrogen;

10

(b) wherein n is 0;


(c) wherein X1, X2, X3, X4, X5, and X6 are independently selected from the group consisting of an hydrido, a chloro, a fluoro, a bromo, and an iodo;

(d) wherein m is 0, 1, 2, or 3; and

(e) wherein the compound is a cytochrome P450 reaction product.

15

4. The compound of claim 1 having **Formula III**:



(a) wherein R1 and R2 are each phenyl; and wherein the phenyl contains an aryl ring carbon and the $(CH_2)_n$ or $(CH_2)_m$ is coupled via a covalent bond to the aryl ring carbon;

20

(b) wherein n is 1;

(c) wherein X1, X2, X3, X4, X5, and X6 are independently selected from the group consisting of a chloro, a fluoro, a bromo, and an iodo;

(d) wherein m is 1; and

(e) wherein the compound is a cytochrome P450 substrate.

5

5. The compound of claim 1, 2, or 3, wherein X1, X2, X3, X4, X5, and X6 are hydrido.

6. The compound of claim 1, wherein the compound is selected from the group of
10 compounds in which:

(a) X2 and X5 are independently selected from the group consisting of chloro and fluoro;

(b) R1 is phenyl and n is 1;

(c) R1 is phenyl, n is 1, and X2 and X5 are independently selected from the group
15 consisting of chloro and fluoro;

(d) R1 is phenyl, R2 is phenyl, n is 1, and m is 1;

(e) R1 is phenyl, R2 is phenyl, n is 1, m is 1, and X2 and X5 are independently selected from the group consisting of chloro and fluoro;

(f) R2 is phenyl and m is 1; and

(g) R2 is phenyl, m is 1, and X2 and X5 are independently selected from the group
20 consisting of chloro and fluoro.

7. The compound of claim 1, wherein the compound has a K_m of greater than about 50 nM with a V_{max} greater than about 0.05 min^{-1} for the cytochrome P450-catalyzed
25 reaction.

8. A composition comprising a compound of any of claims 1-7, wherein the compound is present in the composition at a concentration greater than at least 50% by weight.

30 9. The composition of claim 8, wherein the compound is present in the composition at a concentration greater than at least 75% by weight.

10. The composition of claim 8, wherein the compound is present in the composition at a concentration greater than at least 80% by weight.

5 11. The composition of claim 8, wherein the compound is present in the composition at a concentration greater than at least 90% by weight.

12. The composition of claim 8, wherein the compound is present in the composition at a concentration greater than at least 95% by weight.

10

13. A method for assaying cytochrome P450 enzyme activity comprising:
contacting a cytochrome P450 enzyme with a compound selected from the group consisting of:

- 15 (a) a compound of Formula I as defined by claim 2;
(b) a compound of Formula III as defined by claim 4; and
(c) benzyloxyfluorescein benzyl ester (DBF);

under conditions whereby the cytochrome P450 enzyme catalyzes the conversion of the compound to a fluorescent product.

20 14. The method of claim 13, wherein the cytochrome P450 enzyme is contained in a biological sample.

15 15. The method of claim 13, wherein the sample is selected from the group consisting of a liver sample such as a crude homogenate, partially purified, or purified liver enzyme obtained from a biopsy, a cDNA-expressed cytochrome P450, hepatocytes, and microsomes.

16. A screening method for selecting agents which inhibit cytochrome P450 enzyme activity comprising:
30 contacting a cytochrome P450 enzyme with a compound selected from the group consisting of:

- (a) a compound of Formula I as defined by claim 2;
- (b) a compound of Formula III as defined by claim 4; and
- (c) benzyloxyfluorescein benzyl ester (DBF);

in the presence of a putative cytochrome P450 enzyme inhibitor, and under conditions
5 whereby the cytochrome P450 enzyme catalyzes the conversion of the compound to a
fluorescent product; and

selecting an agent which inhibits cytochrome P450 enzyme activity as a cytochrome
P450 enzyme inhibitor.

10 17. The method of claim 16, wherein the compound is contacted with the putative
cytochrome P450 enzyme inhibitor in a multiwell plate well.

18. A method for visualizing a cytochrome P450 enzyme comprising: contacting a
cytochrome P450 enzyme-containing sample with a compound selected from the group
15 consisting of:

- (a) a compound of Formula I as defined by claim 2;
- (b) a compound of Formula III as defined by claim 4; and
- (c) benzyloxyfluorescein benzyl ester (DBF);

and subjecting the cytochrome P450 enzyme and the compound to conditions whereby
20 the cytochrome P450 enzyme catalyzes the conversion of the compound to a fluorescent
product.

19. The method of claim 18, wherein the cytochrome P450 enzyme-containing
sample is a tissue section.

25 20. A product produced by the process of allowing a cytochrome P450 enzyme to
react with a substrate that is a compound selected from the group consisting of:

- (a) a compound of Formula I as defined by claim 2;
- (b) a compound of Formula III as defined by claim 4; and
- 30 (c) benzyloxyfluorescein benzyl ester (DBF);

under conditions whereby the cytochrome P450 enzyme converts the substrate to a compound of Formula II.

1/5




Fig. 1




Fig. 2

2/5




Fig. 3A






Fig. 3B

3/5



SUPERSOMES


Fig. 3C




SUPERSOMES

Fig. 3D

4/5



SUPERSOMES
Fig. 3E



SUPERSOMES
Fig. 3F

5/5




Fig. 4A




Fig. 4B

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 00/21894

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D311/82 C12Q1/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07D C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 94 05688 A (APPLIED BIOSYSTEMS, INC.) 17 March 1994 (1994-03-17) the whole document	1-20
Y	WO 97 39064 A (MOLECULAR PROBES, INC.) 23 October 1997 (1997-10-23) the whole document	1-20
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *A* document member of the same patent family

Date of the actual completion of the international search

1 February 2001

Date of mailing of the international search report

14/02/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Beslier, L

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 00/21894

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	MACE ET AL: "Development of CYP450-expressing human bronchial epithelial cell lines for in vitro pharmacotoxicologic applications" IN VITRO TOXICOLOGY,US,MARY ANN LIEBERT, NEW YORK, NY, vol. 10, no. 1, 21 March 1997 (1997-03-21), pages 85-92, XP002115754 ISSN: 0888-319X the whole document ---	1-20
Y	HAASCH M L ET AL: "USE OF 7-ALKOXYPHENOXAZONES, 7-ALKOXYCOUMARINS AND 7-ALKOXYQUINOLINES AS FLUORESCENT SUBSTRATES FOR RAINBOW TROUT HEPATIC MICROSOMES AFTER TREATMENT WITH VARIOUS INDUCES" BIOCHEMICAL PHARMACOLOGY,GB,PERGAMON, OXFORD, vol. 47, no. 5, 2 March 1994 (1994-03-02), pages 893-903, XP000901180 ISSN: 0006-2952 the whole document ---	1-20
P,Y	WO 00 35900 A (AURORA BIOSCIENCES CORP.) 22 June 2000 (2000-06-22) particularly tables 12 and 14 ---	1-20
P,Y	WO 99 58710 A (SMITHKLINE BEECHAM PLC) 18 November 1999 (1999-11-18) the whole document ---	1-20
P,Y	WO 00 04008 A (GENTEST CORP.) 27 January 2000 (2000-01-27) cited in the application the whole document ---	1-20
X	TRAN J.: "Intramolecular triplet- triplet energy transfer from xanthene dyes to an anthryl substituent" JOURNAL OF PHOTOCHEMISTRY AND PHOTOBIOLOGY, A: CHEMISTRY., vol. 71, no. 1, 1993, pages 45-49, XP000979202 ELSEVIER SEQUOIA, LAUSANNE., CH ISSN: 1010-6030 the whole document --- -/--	1-7

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 00/21894

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	AMAT-GUERRI F.: "Synthesis and spectroscopic properties of new Rose Bengal and Eosin Y derivatives" DYES AND PIGMENTS., vol. 12, no. 4, 1990, pages 249-272, XP000979190 ELSEVIER APPLIED SCIENCE PUBLISHERS. BARKING., GB ISSN: 0143-7208 the whole document ----	1-7
X	HE J.: "Mechanism of primary photochemical reaction of eosin bis (diphenyliodonium) salt" CHINESE SCIENCE BULLETIN., vol. 41, no. 1, 1996, pages 38-43, XP000979197 SCIENCE PRES, BEIJING., CN the whole document ----	1-7
X	EP 0 353 953 A (ICI PLC) 7 February 1990 (1990-02-07) the whole document ----	1-7
A	WO 87 03541 A (POLAROID CORP.) 18 June 1987 (1987-06-18) page 32, line 1-6 -----	1-7

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/21894

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9405688 A	17-03-1994	DE 69324494 D	20-05-1999
		DE 69324494 T	23-09-1999
		EP 0658167 A	21-06-1995
		JP 8505121 T	04-06-1996
		US 6096723 A	01-08-2000
		US 5654442 A	05-08-1997
		US 5885778 A	23-03-1999
WO 9739064 A	23-10-1997	US 6162931 A	19-12-2000
		AU 717569 B	30-03-2000
		AU 2801297 A	07-11-1997
		CA 2222275 A	23-10-1997
		EP 0853647 A	22-07-1998
		JP 11508277 T	21-07-1999
WO 0035900 A	22-06-2000	US 6143492 A	07-11-2000
		AU 3119000 A	03-07-2000
WO 9958710 A	18-11-1999	NONE	
WO 0004008 A	27-01-2000	US 6130342 A	10-10-2000
EP 353953 A	07-02-1990	AU 3935589 A	17-05-1990
		JP 2163099 A	22-06-1990
		US 5063149 A	05-11-1991
WO 8703541 A	18-06-1987	US 5196297 A	23-03-1993
		CA 1296010 A	18-02-1992
		CA 1316740 A	27-04-1993
		DE 3670803 D	07-06-1990
		EP 0250558 A	07-01-1988
		JP 2628607 B	09-07-1997
		JP 7010869 A	13-01-1995
		JP 6098835 B	07-12-1994
		JP 63501941 T	04-08-1988
		US 4970309 A	13-11-1990
		US 5028725 A	02-07-1991
		US 5220036 A	15-06-1993

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)